## <u>Anecdotal Description of Mosquito Species Occurring in Maine and New Hampshire:</u>

Compiled by Richard Dearborn and Kimberly A. Foss: Maine Department of Conservation, Forest Health and Monitoring, Insect and Disease Lab 2003, from a variety of sources. Revised by Kimberly A. Foss, SWAMP, Inc/Municipal Pest Management Services, Inc. October 2015.

GENUS species	Current Disease Associations (X)=primary vector	Estimated Flight Range	Bites Humans	Adult Host	Larval Habitat	No. Gen./ Yr.	Over- winter Stage	Common Names and Comments
AEDES								
cinereus (Meigen)	WNV EEE SLE	100 to 1000 feet	Yes Major pest in wooded or shaded areas	Mammals	Wooded snowmelt pools, semi-permanent bogs and swamps	2-3	Egg	Day and night biter
vexans (Meigen)	WNV EEE	5 to 10 miles	Yes Major pest	Mammals	Wooded temporary, permanent, semipermanent pools, open flooded areas	2-3	Egg	Day and night biter
OCHLEROTATUS								
abserratus (Felt & Young)			Yes Common spring pest	Mammals, birds	Snowmelt pools	1	Egg	Day and night biter
atropalpus (Coquillett)	WNV	100 to 1000 feet	Yes Around breeding areas	Mammals	Rock pools, some artificial containers	1	Egg	Day and night biter
aurifer (Coquillett)		½ mile	Yes Around breeding areas	Mammals	Snowmelt pools, swamps, bogs, open marshes	1	Egg	Day and night biter
canadensis (Theobald)	WNV EEE	½ mile	Yes Major late spring pest around breeding areas	Mammals, amphibians, reptiles, sometimes birds	Wooded snowmelt pools, flood waters	1-2	Egg	Day and night biter

cantator (Coquillett)	WNV EEE		Yes	Mammals, birds	Salt marshes, fresh or brackish water	1+	Egg	Day and night biter
communis (DeGeer)			Yes	Mammals, birds	Wooded snowmelt pools	1	Egg	Day and evening biter
decticus (Howard, Dyar and Knab)			Yes	Mammals, birds	Sphagnum, acid bogs	1	Egg	Day and night biter
diantaeus (Howard, Dyar and Knab)			Yes Wooded areas	Mammals, birds	Wooded snowmelt pools	1	Egg	Morning and evening biter
dorsalis (Meigen)	WNV SLE	10 to 20 miles	Yes	Large mammals, sometimes large birds	Temporary freshwater and brackish pools marshes and ditches	1+	Egg	"Pale marsh mosquito" New record for 2003 (M. Holman)
excrucians (Walker)		½ mile	Yes Common spring- summer pest	Mammals, sometimes birds	Wooded snowmelt pools, marshes	1-2	Egg	Day and evening biter
fitchii (Felt & Young)	WNV	About 1 mile	Yes Common spring- summer pest in wooded areas	Mammals, birds	Snowmelt pools, bogs, grassy roadside ditches	1	Egg	Day and night biter
hendersoni (Cockerell)		About 1 mile	Yes	Mammals	Tree holes, occasionally tires	1-2	Egg	
implicatus (Vockeroth)			Yes Spring pest	Mammals	Wooded snowmelt pools	1	Egg	Day and night biter
intrudens (Dyar)			Yes Common spring pest	Mammals	Wooded snowmelt pools	1	Egg	Day and night biter

japonicus (Theobald)	WNV SLE		Yes	Mammals, birds	Tires, artificial containers, tree holes, rock pools	2+	Egg	Day biter New Record Portland, Maine: June 26, 2001 (K.Foss)
pionips (Dyar)			Rarely		Snowmelt pools	1	Egg	
provocans (Walker)	WNV		Yes Early spring	Mammals	Semipermenent marshes, wooded snowmelt pools	1	Egg	Evening biter
punctor (Kirby)			Yes Spring	Mammals	Wooded snowmelt pools	1	Egg	Day and night biter
riparius (Dyar and Knab)								New record for 2003 (M. Holman)
sollicitans (Walker)	WNV EEE	100 miles or more	Yes Major coastal summer pest	Mammals, birds, reptiles, amphibians	Salt marshes	4+	Egg	"Eastern salt marsh mosquito", Day and night biter
sticticus (Meigen)	WNV	4 miles	Yes Major pest around breeding areas	Mammals, birds, reptiles	Flood waters, wooded snowmelt pools	1-2	Egg	Day and evening biter
stimulans (Walker)	WNV	2 miles	Yes Major spring pest	Mammals, birds	Snowmelt pools	1	Egg	Long lived
taeniorhynchus (Wiedemann)	WNV EEE		Yes Major pest around breeding areas	Birds, mammals	Salt marshes	2+	Egg	"Black salt marsh mosquito", Day and evening biter New record for 2002(M. Holman)
triseriatus (Say)	WNV LAC (X) EEE	½ to 1 mile	Yes Common summer pest around breeding	Mammals, birds, reptiles, amphibians	Tires, artificial containers, tree holes	1	Egg	"Tree hole mosquito" Day and evening biter

			areas					
trivittatus (Coquillett)	WNV EEE	½ mile	Yes Common summer pest around breeding areas	Mammals, birds	Wooded snowmelt pools, floodwaters	1	Egg	Day and evening biter
ANOPHELES								
barberi (Coquillett)	WNV		Yes	Mammals, sometimes birds	Tree holes, artificial containers	1-2	Larva	New record for 2004 (M. Holman)
crucians	WNV EEE Malaria	<1mile	Yes	Mammals, preferably livestock over humans	Grass fields, retention ponds, detention ponds, and marshes	2+	Larva	New record for NH 2015 (K. Foss)
earlei (Vargas)		1 to 2 miles	Yes Common spring pest	Mammals	Confined bodies of water	1-2	Adult	Day and night biter
punctipennis (Say)	WNV Malaria	1 to 2 miles	Yes Major summer pest	Mammals, birds	Confined and flowing bodies of water, artificial containers	2-3	Adult	"Spotted- winged Mosquito", Day and night biter
quadrimaculatus (Say)	WNV Malaria (X)	1 mile	Yes Common summer pest	Mammals, sometimes birds and reptiles	Confined bodies of water	2-3	Adult	Common "Malaria Mosquito", Day and night biter
walkeri (Theobald)	WNV Malaria	1 to 2 miles	Yes	Mammals	Confined bodies of water	2+	Egg	Day and night biter
COQUILLETTIDIA								
perturbans (Walker)	WNV EEE	1 to 10 miles	Yes Major summer pest	Birds, mammals, amphibians, sometimes reptiles	Cattail marshes	1-2	Larva	Larvae attach to the base of aquatic plants Day and night biter
CULEX								

pipiens (Linnacus)	WNV (X) SLE (X) EEE	1 mile or more	Rarely	Birds, rarely mammals	Artificial containers, grassy roadside ditches, catch basins	1-2	Adult	"Northern house mosquito"
restuans (Theobald)	WNV (X) SLE (X) EEE	1 mile	Yes	Birds, sometimes mammals	Tires, tree holes, artificial containers, puddles, grassy roadside ditches, catch basins	1-2	Adult	Day and night biter
salinarius (Coquillett)	WNV (X) SLE (X) EEE		Yes	Birds, mammals	Artificial containers, grassy roadside ditches, brackish water, catch basins	1-2	Adult	Night biter, enters homes
territans (Walker)	WNV EEE	1 mile	Rarely	Cold blooded vertebrates (e.g. frogs), rarely birds	Pond edges, pools, marshes, grassy roadside ditches, artificial containers	1-3	Adult	
CULISETA								
impatiens (Walker)	WNV		Yes Uncommon early spring species	Mammals	Semipermanent ponds, bogs, wooded ground pools	1	Adult	Long lived, rare, day and evening biter
inornata (Williston)	WNV EEE		Yes Uncommon early spring species	Mammals	Wooded snowmelt pools, marshes, bogs, swamps	2+	Adult	"Winter mosquito"
melanura (Coquillett)	WNV EEE (X)	100 to 1000 yards	Rarely	Birds	Within stumps in acidic swamps and bogs, snowmelt pools	2+	Larva	
minnesotae (Barr)			Rarely	Birds, small mammals, turtles	Snowmelt pools, marshes	1-2	Adult	New Record for 2001 (M. Holman)
morsitans (Coquillett)	WNV EEE		Rarely	Birds	Semipermanent swamps, wooded snowmelt pools, marshes, bogs	1	Egg	

PSOROPHORA								
ciliata (Fabricius)	WNV EEE	5 to 10 miles	Yes, day and night biter Uncommon	Mammals	Temporary open sunlit rain filled fields and flood-water areas	1+	Egg	"Gallinipper" New Record for 2006 S. Berwick (K. Foss)
ferox (Humboldt)	WNV EEE	Up to 1 mile	Yes Within wooded areas, Uncommon species	Mammals	Wooded temporary ground pools, flood-water areas	1	Egg	"White- footed woods mosquito", day and evening biter New Record for 2001 (M. Holman)
URANOTAENIA								
sapphirina (Osten Sacken)	WNV	Up to 8 miles	Rarely Summer species	Birds	Permanent and semipermanent ponds, pools, swamps, marshes	1-2	Adult	New Record Portland, Maine: July 24, 2001 (K. Foss)
WYEOMYIA								
smithii (Coquillett)			Never	Feeds as larvae on other insects in pitcher plant fluid	Sphagnum bogs	1	Larva	"Pitcher plant mosquito" spends most of the year in larval stage
ORTHOPODOMYIA							_	
signifera (Coquillett)	WNV EEE	Less than 100 ft	Rarely, uncommon species slow to develop	Birds	Deep tree rot holes and wooden containers	2+	Egg in north, larvae in south	